

Antihelminth (Worm) Treatment

Albendazole 400mg Tablets			
	Children 12-24 Months	All others	
Routine treatment in endemic areas	½ tablet single dose	1 tablet single dose	
Recommended every 3 months			
Treatment of heavy worm burden OR	½ tablet twice daily for	1 tablet twice daily for	
Failed single dose treatment	3 days	3 days	

Mebendazole 100mg Tablets		
*Not Recommended For Children Under 2 Years of Age		
	All persons over 2 years old	
Routine treatment in endemic areas	1 tablet single dose	
Recommended every 3 months		
Treatment of heavy worm burden OR	1 tablet twice daily for 3 days	
Failed single dose treatment		
Inadequate response to treatment of heavy worm	1 tablet twice daily for 3 days	
burden	2 weeks after first course	

Mebendazole 500mg Tablets		
*Not Recommended For Children Under 2 Years of Age		
(500mg single use recommended for suspected non-	All persons over 2 years old	
compliance)		
Treatment of heavy worm burden OR	1 tablet single dose	
Failed single dose treatment		
Inadequate response to treatment of heavy worm	1 tablet single dose 2 weeks	
burden	after first course	

lvermectin		
	All Patients	
Treatment of heavy worm burden OR	0.2mg/kg as a single dose. May	
Failed single dose treatment	repeat in 2 weeks if necessary	

Additional Information:

Albendazole and Mebendazole are chewable tablets and can be administered immediately in the clinic while the patient is observed taking the medication.

Use in pregnancy: Mebendazole and Albendazole are Pregnancy Category C and not recommended for long term use. Ivermectin is contraindicated in pregnancy.

Iron supplements and vitamins may be needed to treat related co-morbidities such as anemia and malnutrition.